



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTES UPON VARIATION OF *MELISSODES LUPINA* CRESSON.

By T. D. A. COCKERELL, Las Vegas, N. M.

THREE male specimens of *Melissodes* collected by E. S. Tucker at Colorado Springs, Colo., during August, 1906, were submitted to me for examination.

The examples are so close to *Melissodes lupina* Cr. that I doubt whether they are more than a geographical race. Comparing them with a cotype of *M. lupina*, I find that the eyes of these specimens are bluer green and diverge less above, and the mesothorax is more shiny. A specimen from Raton, N. M., which I had labeled *lupina*, has the same color of the eyes and shiny mesothorax as in Mr. Tucker's insects, but the eyes diverge as in typical *lupina*. The color of the eyes is not quite satisfactory when examined only in dry specimens. On the whole, I should call these bees *M. lupina*, but they might be regarded as a distinct subspecies. Were it not for the Raton specimen being in some degree intermediate, I should not feel so undecided.

Mr. Tucker's insects, compared with cotype of *M. aurigenia*, have the fourth antennal joint shorter and the pubescence different in color. Compared with cotype of *M. agilis*, these specimens have the face narrower (which may vary) and the apical plate of abdomen conspicuously broader.